



Bridging the Protein Gap: Navigating Variable Milk Composition and Delivery in High-Risk Neonates and Infants



ANNENBERG CENTER FOR HEALTH SCIENCES
AT EISENHOWER
Imparting knowledge. Improving patient care.

This activity is supported by an educational grant from Mead Johnson Nutrition.



Faculty Presenters

Sarah Fleet, MD, PNS

Program Director, Advanced Nutrition
Training Program

Pediatric Gastroenterology, Hepatology
& Nutrition

Boston Children's Hospital

Assistant Professor of Pediatrics

Harvard Medical School

Boston, Massachusetts

Ting Ting Fu, MD, MS

Division of Neonatology

UC Department of Pediatrics

Cincinnati Children's Hospital
Medical Center

Cincinnati, Ohio



Faculty Disclosures

It is the policy of the Annenberg Center to ensure fair balance, independence, objectivity, and scientific rigor in all programming. All faculty participating in accredited programs are expected to identify and reference off-label product use and disclose any relationship with those supporting the activity or any others whose products or services are discussed.

Sarah Fleet, MD, PNS

No relationships to disclose.

Ting Ting, Fu, MD, MS

Speakers Bureau

Medela

Faculty have documented that this presentation will involve discussion of unapproved or off-label, experimental, or investigational use. For complete disclosure information, please see our course syllabus.




Learning Objectives


By participating in this education, you will better:



Describe the differences in nutrient composition and bioactive components in mother's own milk vs donor human milk



Identify the impact of cumulative protein deficits on clinical and growth outcomes in both preterm infants and term infants with medical complexity



Analyze the latest clinical guidelines for fortifying human milk to meet the heightened metabolic demands of the medically complex neonatal and pediatric populations





Introduction: Why the Emphasis on Protein?

Ting Ting Fu, MD, MS




Amino Acids and Protein During Fetal Development

Amino acids serve critical functions during fetal development.

- Energy
- Growth and organ development
- Cell structure and maturation
- Cellular remodeling
- Immune system development




Protein Accretion and Body Composition by Gestational Age in the Human Fetus



The placenta supplies high levels of amino acids to the fetus to meet the energy and nutritional requirements for rapid fetal growth, with most protein deposition occurring in the third trimester.



Protein Accretion and Body Composition by Gestational Age in the Human Fetus



Preterm birth abruptly interrupts critical periods of in utero protein delivery.



Goals for Preterm Infant Growth

2022 ESPGHAN Growth Targets for Preterm Infants¹:

- Regain birth weight by days 7 to 10 after initial loss of up to 10% at days 3 to 4
- After days 7 to 10, achieve **growth along a target fetal growth percentile**, with gradual transition to corresponding percentile on postnatal growth references
- Promote catch-up growth in infants with growth faltering while avoiding very rapid catch-up growth

Adequate protein intake is key for meeting growth targets.¹

Fenton Fetal-Infant Growth Chart for Preterm Girls²






Image reprinted under a Creative Commons license. ©Fenton TR, Kim JH. *BMC Pediatr.* 2013;13:59. (CC BY 2.0).



Actual Target Velocities for Preterm Infants




Enteral Protein Intake and Growth in Preterm Infants

Systematic Review and Meta-Analysis

- 10 RCTs comparing higher vs lower enteral protein intake
- Included 646 preterm infants born at <32 weeks GA
- Analyses adjusted for concurrent energy intake

↑ 8.8 change in weight gain for each g/kg/day increase in protein intake ($P \leq .001$)
g/kg/day

Association Between Protein Intake and Weight Gain



Protein Requirements in Infancy and Childhood

Protein requirements are highest for VLBW and preterm infants and decline with older age.¹

Recommendations for Protein Intake¹⁻³

VLBW/Preterm Infant

3.5-4.0
g/kg/day*

*Up to 4.5 g/kg/day in
infants not meeting
growth targets

Term Infant

2.0-2.5
g/kg/day

Older Infant

1.5-2.0
g/kg/day

Toddler

1.0-1.5
g/kg/day

Adolescent

0.8-1.5
g/kg/day

Note: During periods of critical illness and catch-up growth, protein requirements may be up to 20% to 50% higher!

Protein Requirements in Infancy and Childhood

Protein requirements are highest for VLBW and preterm infants and decline with older age.¹

Recommendations for Protein Intake¹⁻³

VLBW/Preterm Infant

3.5-4.0
g/kg/day*

*Up to 4.5 g/kg/day in infants not meeting growth targets

Term Infant

2.0-2.5
g/kg/day

Older Infant

1.5-2.0
g/kg/day

Toddler

1.0-1.5
g/kg/day

Adolescent

0.8-1.5
g/kg/day


Note: During periods of critical illness and catch-up growth, protein requirements may be up to 20% to 50% higher!



Protein Requirements in Infancy and Childhood

Protein requirements are highest for VLBW and preterm infants and decline with older age.¹

Recommendations for Protein Intake¹⁻³




1. Herrera OR, Helms RA. In: Corkins MR et al (eds). *ASPEN Pediatric Nutrition Support Core Curriculum, 3rd Edition*. American Society for Parenteral Nutrition (ASPEN);2025. 2. Koletzko B et al, eds. *Nutritional Care of Preterm Infants. Scientific Basis and Practical Guidelines, 2nd ed.* Karger; 2021. 3. Embleton ND et al. *J Pediatr Gastroenterol Nutr.* 2023;76(2):248-268.



Human Milk Composition and the Neonatal Protein Gap

Human Milk Composition¹



⚠ Think of “g/dL” as %


Milk source	Protein concentration	Protein intake at recommended preterm feeding volumes	
		140 mL/kg/day ⁴	200 mL/kg/day ⁴
Preterm milk (MOM)	1.6-2.2 g/dL ²	~2.3 g/kg/day	~3.2 g/kg/day
Term milk (DHM)	0.9-1.2 g/dL ³	~1.3 g/kg/day	~1.8 g/kg/day

Unfortified human milk is associated with a **protein gap of 0.3-3.3 g/kg/day** when using the recommended protein intake of **3.5-4.5 g/kg/day**.

1. Perrin MT et al. *Adv Nutr.* 2020;11(4):960-970. 2. Gates A et al. *Am J Clin Nutr.* 2021;114(5):1719-1728. 3. Kim SY, Yi DY. *Clin Exp Pediatr.* 2020;63(8):301-309. 4. Koletzko B et al, eds. *Nutritional Care of Preterm Infants. Scientific Basis and Practical Guidelines, 2nd ed.* Karger; 2021.



Daily Gaps in Protein Intake Can Yield Cumulative Deficits Over Time






NICU: Human Milk Composition and the Protein Gap

Ting Ting Fu, MD, MS



Not All Human Milk Is Equal (or 20 kcal/oz)



Colostrum → Transitional → Mature



3 days

5 days

6 days

25 days

Foremilk vs Hindmilk



Top right image reprinted under a Creative Commons license. ©Amada44. https://commons.wikimedia.org/wiki/File:From_Colostrum_to_Breastmilk_-_4241.jpg. (CC BY-SA 3.0). Bottom right image unpublished from Fu TT.




Preterm Human Milk Composition: Declining Protein Concentration Later in Lactation

Prospective, Longitudinal Cohort Study

- Women who delivered ≤ 33 weeks' gestation (N = 38)
 - Average GA of 28.2 weeks, with 42% delivering < 28 weeks
- Pooled 24-hour milk samples
- Assessed macro- and micronutrient composition

Protein Concentration by Lactation Stage




Consistently Lower Protein Levels in DHM

Observational Study

- Analyzed DHM samples (n = 15) from 7 sources
- Compared nutrient profiles of DHM to those of:
 - Early preterm human milk (day 7; n = 36)
 - Mature preterm human milk (day 28; n = 28)

Protein Concentration in DHM and Preterm Milk



* $P < .05$; **** $P < .0001$



Nutrient Variability in DHM: Potential Sources

In a systematic review and meta-analysis, the macronutrient composition of DHM varied by ≥ 2 fold

Potential Sources of DHM Variability



Donors

- Gestation stage (term vs preterm)
- Lactation stage
- Maternal diet
- Collection method
- Collection time



Milk Banks

- Pasteurization processes
- Use of pre-pooling analysis
- Mixing practices

Note: Pooling DHM can reduce variability and is an important consideration for milk banks.



Key Characteristics of DHM Impacting Composition



Heat-pasteurized to destroy microbes, which can also reduce activity and levels of bioactive components



Primarily expressed by healthy mothers of term infants at later lactation stages



Pooled from multiple mothers to reduce donor-to-donor and lot-to-lot variability



Donor Milk Pasteurization: Effects on Milk Composition

Changes in bioactive components¹⁻³

- Complete loss of certain **enzymes** and maternal **cell populations** (eg, neutrophils, stem cells)
- Reduced activity level or concentration of other **enzymes, cytokines, growth factors, immunoglobulins, and hormones**

Changes in micronutrient composition¹

- Reduced **ascorbic acid** and **vitamin B6**

 Human Milk Banking Association of North America (HMBANA) uses Holder pasteurization (62.5 °C for 30 minutes), but other sources may use different pasteurization methods.⁴



Measuring Protein Levels in Human Milk

Gross protein is the total nitrogen content and includes^{1,2}:

- **True protein** (nitrogen in amino acids), the metabolizable content providing necessary nutrients for growth and maintenance
- **Non-protein nitrogen (NPN)** (nitrogen in compounds such as urea and ammonia)



Total Protein = True Protein + NPN

(NPN estimated to be ~20-30% of N in human milk)²



Measuring Protein Levels in Human Milk

Total Protein
(total nitrogen content)

=

True Protein

- Nitrogen from amino acids
- Provides nutrients for growth and maintenance

+

Non-Protein Nitrogen (NPN)


- Non-nutritional nitrogen compounds (eg, urea, ammonia)
- Estimated to be ~20%-30% of nitrogen in human milk)²



DHM Labels: What's On Them & What's Not?

PASTEURIZED HUMAN MILK

Local Milk Bank, Inc.



Batch: 012345-01

Milk type: Term

Exp: JAN 01, 2027

Volume: 100 mL

Cal/oz: 20 Prot: 1.0 g/dL

May indicate milk type (eg, term, preterm, targeted)

Doesn't distinguish total vs true protein

Nutritional information dependent on milk bank having a human milk analyzer (HMA)



Not included on labels:

- Information on carbohydrate or fat levels (in most cases, may be included on the packing sheet)
- Micronutrients
- Bioactive components
- Pooling method or number of donors pooled



Accuracy of Donor Milk Labels¹⁻³

- Secondary analysis of **101 donor milk samples** from an ongoing prospective cohort study³
 - Rigorous sample collection with mid-infrared HMA analysis in triplicate
- Calorie and protein levels **not** statistically different from labeled values
- Intra-batch variability (within the same pool) can occur due to mixing and bottling practices if not using automated processes




Overestimation of Protein in DHM

Observational Study


- Triplicate measures of total nitrogen, NPN, and amino acid profiles from DHM samples
- Values compared against labeled and calculated protein content (using assumption of 20% NPN)

Overestimation of Reported Protein Content on DHM Labels (n = 6)



0.15 g/dL median overreporting of protein content on DHM labels

Overestimation of Calculated Total Nitrogen in DHM (n = 15)




33% mean NPN content in DHM



Key Takeaways: the Protein Gap

- Human milk alone does **not** meet recommended protein targets for preterm infants
- Human milk composition is variable, with differences in protein levels by preterm vs DHM and other factors
- Nutritional **interventions are needed to meet protein requirements** to optimize growth and development in preterm infants





NICU: Bridging the Gap With Fortification

Ting Ting Fu, MD, MS



Methods of Human Milk Fortification

Standard

most common & easiest

- Fixed amount of fortifier added to fixed milk volume
- Based on manufacturer's instructions (typically assumes starting protein and energy content of 1.5 g/dL and 20 kcal/oz, respectively)

Adjustable

more cost efficient & less labor intensive than targeted

- Protein concentration adjusted based on serial blood urea nitrogen (BUN) measurements
- Additional protein supplementation added if level is <10 mg/dL

Targeted

most accurate & most costly

- Macronutrient concentrations in human milk analyzed with an HMA
- Fortification procedures based on analysis



Standard Fortification Using Traditional Assumptions May Not Be Sufficient

Standard¹

most common & easiest




- Fixed amount of fortifier added to fixed human milk volume
 - Based on manufacturer's instructions, which typically assumes starting protein and energy content of 1.5 g/dL and 20 kcal/oz, respectively
- Traditional assumptions regarding human milk composition have recently been challenged^{1,2}
 - Only 11% of human milk samples fortified using standard methods reached 4 g/kg/day protein²
 - **No** samples reached 4.5 g/kg/day protein with standard fortification



Comparison of Standard and Targeted Fortification

Double-Blind Randomized Controlled Trial

- Enrolled infants <30 weeks GA
- Compared standard fortification (n = 51) with standard fortification plus targeted fortification using modular protein, lipids, and carbohydrates (n = 52)
- Analyzed milk samples (n = 2810) and infant body composition



* $P < .001$

Middle image used under a Creative Commons license. ©Rochow N et al. *Clin Nutr.* 2021;40(1):54-63. (CC BY-NC-ND 4.0).



Baseline Protein Enrichment of DHM

- Initial **prospective cohort study** of 69 VLBW infants who received **target-pooled donor milk (20 kcal/oz)**¹
 - Protein content was still 0.9 g/dL
 - Change in z scores from birth to 36 weeks PMA:
 - » Weight z score decreased by 0.5
 - » Length z score decreased by 1.0
- A **protein-enriched donor milk (DBM+)** for clinical use was developed by adding a liquid protein product²
 - 6 mL liquid protein per 90 mL DHM
 - Fed as early as initiation of enteral feeding

Comparison of Protein Levels With DHM and DBM+

	Protein (g/dL) (baseline)	Protein (g/dL) (fortified to 24 kcal/oz)
DHM (20 kcal/oz)	0.9	2.4
DBM+	1.9	3.2

PMA, postmenstrual age



Baseline Protein Enrichment: Growth With MOM, DHM, or DBM+


Prospective Cohort Study

- Compared VLBW infants receiving DHM, DBM+, or MOM as primary feeds
- Evaluated safety and growth
- Limited by lack of randomization, exclusion of certain infants by clinical team due to size or illness, and inadequately powered to detect differences in length trajectory

Conclusions

- Appears to be safe in stable VLBW infants
- No difference in incidences of NEC or feeding intolerance
- Weight gain greatest with MOM
- Linear growth with DBM+ comparable to MOM

Comparison of Growth by Primary Feed Type



Enhanced Fortification: MAGIC Study



Not exclusive to
DHM feedings

Ongoing Prospective Cohort Study

- VLBW infants <33 weeks GA from 2 level III NICUs
- Fortified to 26 kcal/oz for **all** VLBW infants after reaching full enteral feeds (day ~12-14)
 - 15 mL bovine milk-based fortifier with extensively hydrolyzed protein per 50 mL human milk^a
- Representative samples of human milk fed to infants collected and analyzed
 - MOM: 24-hour pools sampled up to 3 times per week, depending on supply (n = 198)
 - DHM: 1 bottle from each lot sampled (n = 168)


Enrollment Criteria

Inclusion	Exclusion
<ul style="list-style-type: none">• Birth weight <1500 grams• GA at birth <33 completed weeks• Parental clinical consent to the provision of DHM• Attainment of full enteral feeding volume with fortification to 26 kcal/oz within the first 30 days of life, per clinical team	<ul style="list-style-type: none">• Significant chromosomal/genetic abnormalities impacting growth potential


a. Off-label recipe not studied by the manufacturer, but this recipe has been utilized by our NICU since 2016 and is used across Greater Cincinnati NICUs.



Enhanced Fortification: MAGIC Study

 Not exclusive to DHM feedings

Frequency at Which Enhanced Fortification Meets Recommended Protein Targets, Assuming Intake of 150 mL/kg/day



- Lowest targets consistently met with fortification
- Standard fortification did not adequately meet protein targets with DHM
- **Limitations:** results may differ by fluid intake or fortifier; reflects pooling practices from our regional milk bank; infant outcomes pending

a. Off-label recipe not studied by the manufacturer, but this recipe has been utilized by our NICU since 2016 and is used across Greater Cincinnati NICUs. Image reprinted under a Creative Commons license. © Fu TT et al. *Pediatr Res*. Published online ahead of print January 15, 2026. (CC BY).



Fortification Timing: Beginning Fortification at Onset of Feeding



Not exclusive to
DHM feedings

Retrospective Study¹

- Infants <31 weeks GA (n = 95)
- Compared early fortification at time of first feed vs delayed fortification at 50-100 mL/kg/day using bovine milk-based powdered fortifier

Conclusions

- No difference in weight gain
- No difference in incidence of NEC

Randomized Controlled Trial²

- Infants <29 weeks GA (n = 150)
- Randomized to receive human milk-based fortifier at onset of feeding or a standard unfortified diet prior to usual bovine milk-based fortifier at 14 days

Conclusions

- Improved length and head circumference with early fortification
- No difference in fat-free mass accretion
- No difference in incidence of NEC or spontaneous intestinal perforation



Increased Volumes of Human Milk or Formula May Increase Growth



Not exclusive to
DHM feedings

Randomized Clinical Trial

- Enrolled infants born at <32 weeks' gestation weighing 1001-2500 g
- After reaching full enteral feeds, randomized infants to receive:
 - Higher-volume feeds (180-200 mL/kg/d)
 - Usual-volume feeds (140-160 mL/kg/d)
- Included infants fed fortified human milk and preterm formula

	Higher volume (n = 104)	Usual volume (n = 113)	P value
Mean growth velocity, g/kg/day <i>Primary end point</i>	20.5	17.9	<.001
Mean weight, g	2365	2200	<.001
Head circumference, cm	31.9	31.4	.01
Length, cm	44.9	44.4	.04
Mean (range) days on respiratory support	6 (0-85)	6 (0-85)	0.81
NEC stage ≥2, n (%)	0 (0%)	0 (0%)	1.00
Feeding intolerance, n (%)	2 (2)	3 (3)	1.00



Safety Considerations With High-Protein Fortification

Systematic Review and Meta-Analysis

- Data from 44 randomized and quasi-randomized clinical trials
- Included trials providing protein (both **parenteral** and **enteral**) to preterm infants <37 weeks GA (n = 5338)
- Evaluated long-term effects in children

RISK RATIO
0.95
(95% CI, 0.90-1.01)

Slightly **decreased** chance of **neurodisability-free survival** at age ≥ 12 months ($P = .13$; low-certainty evidence)

RISK RATIO
1.36
(95% CI, 0.89-2.09)

Slightly **increased** chance of **cognitive impairment** at toddler age ($P = .16$; low-certainty evidence)

ADDITIONAL FINDINGS

- **Increased weight** and **head circumference** z scores at discharge
- **No significant differences** in NEC, late-onset sepsis, or other neonatal morbidity markers



Safety Considerations With High-Protein Fortification: Important Study Limitations

Limitations identified in commentaries on the meta-analysis by Das et al:

- Low level of certainty for most findings
- Included studies evaluating parenteral amino acids and enteral protein, which are not comparable interventions
 - Lower protein requirements with parenteral nutrition (PN) vs enteral nutrition
 - Amino acids in parenteral nutrition not well studied
- Lack of adequately powered studies of nutrition intake in preterm infants
- Need for a more holistic approach focused on all macronutrients



Safety of Protein Fortification

- Routine fortification to **protein levels of 3.5-4.0 g/kg/day** is safe and beneficial for growth
- Consider increasing fortification **up to 4.5 g/kg/day** on an individualized basis, including:
 - For infants not meeting growth targets
 - Extremely preterm or VLBW infants
 - Infants on exclusive DHM diets, where standard assumptions may magnify the protein gap
 - Surgical neonates (eg, post-NEC, gastroschisis)




Clinical Integration & Key Takeaways



In concert with individualized nutrition and growth goals, with monitoring

- **Be knowledgeable** about the donor milk product you are selecting and purchasing
 - Understand the processing methods and the limitations
- Although **labels aren't perfect**, if available, they may still be useful to identify a pool to meet the needs of an infant with growth faltering
 - Look for pools with higher protein or energy
- **Prioritize MOM** when possible, and **tailor fortification** strategies for DHM
 - Targeted fortification: ideal, but may not always be practical
 - Universal approach: acknowledge the macronutrient content is low (especially protein) and give more





Protein Metabolism and Requirements in Critically Ill Infants and Children

Sarah Fleet, MD, PNS



Protein Requirements in Infancy and Childhood

Protein requirements are highest for VLBW and preterm infants and decline with older age.¹

Recommendations for Protein Intake¹⁻³

VLBW/Preterm Infant

3.5-4.0
g/kg/day*

*Up to 4.5 g/kg/day in infants not meeting growth targets

Term Infant

2.0-2.5
g/kg/day

Older Infant

1.5-2.0
g/kg/day

Toddler

1.0-1.5
g/kg/day

Adolescent

0.8-1.5
g/kg/day

Note: During periods of critical illness and catch-up growth, protein requirements may be up to 20% to 50% higher¹.


1. Herrera OR, Helms RA. In: Corkins MR et al (eds). *ASPEN Pediatric Nutrition Support Core Curriculum, 3rd Edition*. American Society for Parenteral Nutrition (ASPEN);2025. 2. Koletzko B et al, eds. *Nutritional Care of Preterm Infants. Scientific Basis and Practical Guidelines, 2nd ed.* Karger; 2021. 3. Embleton ND et al. *J Pediatr Gastroenterol Nutr.* 2023;76(2):248-268.



Protein Requirements in Infancy and Childhood

Protein requirements are highest for VLBW and preterm infants and decline with older age.¹


Recommendations for Protein Intake¹⁻³



1. Herrera OR, Helms RA. In: Corkins MR et al (eds). *ASPEN Pediatric Nutrition Support Core Curriculum, 3rd Edition*. American Society for Parenteral Nutrition (ASPEN);2025. 2. Koletzko B et al, eds. *Nutritional Care of Preterm Infants. Scientific Basis and Practical Guidelines, 2nd ed.* Karger; 2021. 3. Embleton ND et al. *J Pediatr Gastroenterol Nutr.* 2023;76(2):248-268.




Metabolism During the Phases of Critical Illness^{1,2}



Catabolism Replaces Energy Loss But Can Have Adverse Outcomes in Children


Metabolic Changes During the Acute Phase of Critical Illness



1. Briassoulis G et al. *Nutrients*. 2024;16(20):3523. 2. Mehta NM. Accessed March 27, 2026. <https://aneskey.com/nutrient-metabolism-and-nutrition-therapy-during-critical-illness/>.



Protein Catabolism in Critical Illness




Muscle loss increases the risk of life-threatening complications:

- Difficulty weaning from ventilation
- Swallowing difficulty

Image reprinted under a Creative Commons license. ©Wilson B, Typpo K. *Front Pediatr.* 2016;4:108. (CC BY).



Goals of Nutritional Support in the PICU




Landmark Study: Adequate Protein Intake Decreases Mortality in the PICU

Prospective Cohort Study

- Included 1245 children from 59 multinational PICUs
- Enrolled children aged 1 month to 18 years who were mechanically ventilated for at least 2 days
- Evaluated mortality based on the percentage of the prescribed daily goal of energy and protein delivered

Mortality by Protein Adequacy and Severity of Illness at Admission



↓ **86%**


decreased odds of 60-day mortality among patients receiving $\geq 60\%$ prescribed protein vs those receiving $< 20\%$ of prescribed protein



Calculating Protein Delivery Targets in the PICU



Do **not** use RDA values



Use **indirect calorimetry** to estimate needs



Minimum protein intake of 1.5 g/kg/day to achieve positive nitrogen balance and avoid cumulative deficits

- Intake of **2.5-3 g/kg/day recommended** for infants and young children on mechanical ventilation



In infants, consider implications of **human milk-based diets** and **potential protein gaps**



PICU Enteral Feeding Considerations: Formula

- Carefully **evaluate infant** and **pediatric formulas** for protein adequacy
 - ~1.4 g/dL protein in most formulas and will **not** meet protein goals for PICU patients^{1,2}
- **To meet protein goals in the PICU, consider^{2,3}:**
 - Adding modular protein supplementation to standard formulas
 - Using high-protein formulas (if tolerated)
- For patients with fluid restrictions or malnutrition, consider **energy- and nutrient-dense formula** feeding, with potential protein supplementation as-needed^{2,3}





Considerations for High-Risk Populations

Sarah Fleet, MD, PNS



Congenital Heart Disease (CHD): High Burden of Malnutrition

- Prevalence of malnutrition ranges from **15%** to **64%**
- Attributed to:
 - Increased cardiac metabolic demands
 - Volume and/or pressure overload
 - Chronic hypoxemia
- **Risk factors for malnutrition:** low birth weight, preterm birth, pulmonary hypertension, pneumonia, congestive heart failure, and age <1 year

Adverse Outcomes of Malnutrition in CHD

- Longer hospitalization
- Greater infection risk
- Higher mortality risk
- Higher risk of adverse neurodevelopmental outcomes
- Worse quality of life



Proposed Feeding Strategies for Infants With CHD

General approach of maximizing nutrition while preventing NEC and fluid overload by using a risk-stratified approach that focuses on protein-dense trophic feeds

Evaluate feeding complication risk based on:

- Demographic risk factors
- Type and severity of CHD

Low risk

Follow protocols for preterm or term infants, allowing term infants to ad lib feed and breastfeed

Monitor feeding tolerance and clinical status

High risk

Initiate trophic feeds at of human milk (if stable) and place access for PN; allow oral feeds with human milk

Additional risk factors (eg, imminent surgery)

Continue trophic feeds and concentrated TPN

No additional risk factors

Advance feeds and monitor tolerance and clinical status



Benefits of Pre- and Perioperative Human Milk Feeds in CHD

Registry-Based Study¹

- Included infants with single ventricle CHD (n = 2491)
- Evaluated human milk-fed infants during the first 2 surgical stages based on feed exposures

Conclusions: Benefits of Human Milk Feeds

- Before stage 1 surgery, decreased risk of preoperative NEC and length of stay
- During stage 1 hospitalization, decreased risk of postoperative NEC, risk of sepsis, and length of stay
- During stage 2 surgery, decreased length of stay

Retrospective Study²

- Included infants with CHD and isolated cardiac lesion at high risk for NEC from single site (n = 546)
- Evaluated preoperative human milk feeding and effects on NEC

Conclusions: Benefits of Human Milk Feeds

- Reduced risk of preoperative NEC after controlling for cardiac lesion, race, feeding volume, birth weight, and preterm birth



Other Complex PICU Populations: Acute Kidney Injury (AKI)

- AKI is a commonly cited reason for inadequate protein intake
- Nitrogen loss from continuous renal replacement therapy (CRRT) is approximately **20% of intake**
- No evidence that higher protein intake in children with AKI is associated with delayed kidney recovery

Nutritional Considerations

- Account for CRRT nitrogen losses when prescribing protein
- Prioritize EN feeding



Other Complex PICU Populations: Postoperative Care

- Nutritional support begins **prior to surgery**:
 - Identify and address malnutrition preoperatively, when possible
 - Avoid prolonged fasting prior to surgery
- After surgery, nutrition support is intended to **restore normal GI function** to allow for nutrient intake and absorption:
 - Sufficient protein intake is required to reduce the impact of catabolism and promote wound healing
 - Early enteral nutrition (within 24 hours) can decrease the risk of nutritional deterioration following surgery



Other Complex PICU Populations: Severe Malnutrition at Risk for Refeeding Syndrome

- Often defined as **weight-for-age or BMI-for-age z score of -3 or less**
 - **Other at-risk patients:** weight loss $\geq 15\%$ in last 3-6 months, little or no nutritional intake for ≥ 5 days, or baseline electrolyte depletion
- Initial hypocaloric feeds should **include adequate protein** (up to goal range)

Initiate hypocaloric feeds (~50%-80% of calories for current weight) with adequate protein and electrolyte supplementation



After ~4 days of stable labs, advance feeds by 10%-20%



Continue advancing calories to meet recommended amount for catch-up growth, with continued protein supplementation



Clinical Integration & Key Takeaways

- **Define and individualize protein targets** in the PICU
- **Incorporate nutritionists** into PICU rounds to support optimal nutrition
- **Aggressively target protein goals** using PN and/or trophic feeds, as needed
- When possible, **assess adequacy** of protein delivery and proactively adjust when indicated
- For infants, **identify milk sources** (DHM vs MOM) and utilize updated reference values when estimating protein content

